

FIELD LIST OF RESERVA COSTANERA SUR - BIRDS RECORDED (May 2014-April 2015)

Cora Rimoldi, J. Simón Tagtachian and Germán Pugnali - www.reservacostanera.com.ar

Anhimidae	Phoenicopteridae	Accipitridae (cont.)	Rallidae (cont.)	Columbidae (cont.)	Picidae (cont.)
<b>F</b> Southern Screamer VP <i>Chauna torquata</i> R	<b>F</b> Chilean Flamingo C <i>Phoenicopus chilensis</i> RV	<b>III</b> Snail Kite L-VP <i>Rostrhamus sociabilis</i> R	<b>II</b> White-winged Coot L-VP <i>Fulica leucoptera</i> R	<b>I</b> Eared Dove B-T <i>Zenaida auriculata</i> R	<b>I</b> Green-barred Woodpecker B-P-T <i>Colaptes melanochloros</i> R
<b>II</b> Fulvous Whistling-Duck L <i>Dendrocygna bicolor</i> R	<b>III</b> Maguari Stork EA-VP <i>Ciconia maguari</i> R	<b>III</b> Long-winged Harrier EA-VP <i>Circus buffoni</i> R	<b>II</b> Sharp-shinned Hawk EA-B <i>Accipiter striatus</i> R	<b>I</b> White-tipped Dove B <i>Leptotila verreauxi</i> R	<b>II</b> Campo Flicker B-T <i>Colaptes campestris</i> R
<b>I</b> White-faced Whistling-Duck L <i>Dendrocygna viduata</i> R	<b>F</b> Wood Stork EA-VP <i>Mycteria americana</i> R	<b>I</b> Roadside Hawk EA-B <i>Rupornis magnirostris</i> R	<b>I</b> Southern Lapwing VP <i>Vanellus chilensis</i> R	<b>F</b> Ash-coloured Cuckoo B <i>Coccyzus cinerea</i> MA	<b>I</b> Southern Caracara EA-B-P <i>Caracara plancus</i> R
<b>F</b> Black-headed Swan L <i>Cygnus melancoryphus</i> R	<b>I</b> Neotropic Cormorant C <i>Phalacrocorax brasilianus</i> R	<b>I</b> Harris's Hawk EA-B <i>Parabuteo unicinctus</i> R	<b>F</b> Black-necked Stilt L <i>Himantopus mexicanus</i> R	<b>III</b> dark-billed Cuckoo B <i>Coccyzus melancoryphus</i> MA	<b>I</b> Chimango Caracara EA-B-P <i>Milvago chimango</i> R
<b>I</b> Coscoroba Swan L <i>Coscoroba coscoroba</i> R	<b>I</b> Ringed Teal L <i>Callonetta leucophrys</i> R	<b>F</b> White-tailed Hawk EA <i>Buteo albicaudatus</i> R	<b>F</b> South American Snipe L <i>Gallinago paraguaiæ</i> MA	<b>I</b> Guira Cuckoo A-B-P-T <i>Guira guira</i> R	<b>III</b> American Kestrel EA <i>Falco sparverius</i> R
<b>I</b> Brazilian Teal L <i>Amazonetta brasiliensis</i> R	<b>I</b> Rufescent Tiger-Heron L-VP-C <i>Tigrisoma lineatum</i> R	<b>F</b> Black-chested B-Eagle EA <i>Geranoaetus melanoleucus</i> R	<b>F</b> Swainson's Hawk EA <i>Buteo swainsoni</i> MB	<b>F</b> Striped Cuckoo A-B-P-T <i>Tapera naevia</i> MA	<b>III</b> Peregrine Falcon EA <i>Falco peregrinus</i> RV
<b>I</b> Yellow-billed Teal L <i>Anas flavirostris</i> R	<b>III</b> Stripe-backed Bittern VP <i>Ixobrychus involucris</i> MA	<b>I</b> Limpkin L-VP <i>Aramus guaranauna</i> R	<b>I</b> Wattled Jacana VP <i>Jacana jacana</i> R	<b>F</b> Tropical Screech-Owl A-B <i>Megascops choliba</i> R	<b>I</b> Monk Parakeet EA-B-A <i>Myiopsitta monachus</i> R
<b>F</b> Yellow-billed Pintail L <i>Anas georgica</i> R	<b>III</b> Black-crowned Night-Heron L-VP <i>Nycticorax nycticorax</i> R	<b>III</b> Striped Heron L-VP <i>Butorides striata</i> R	<b>F</b> South American P-Snipe VP <i>Nycticryphes semicollaris</i> R	<b>III</b> Ferruginous Pygmy-Owl A-B <i>Glaucidium brasilianum</i> R	<b>III</b> Yellow-chevroned Parakeet EA-B <i>Brotogeris chiriri</i> FDE
<b>I</b> Silver Teal L <i>Anas versicolor</i> R	<b>III</b> Cinnamon Teal L <i>Anas georgica</i> R	<b>II</b> Giant Wood-Rail VP-T <i>Aramides ypecaha</i> R	<b>F</b> Limpkin L-VP <i>Aramus guaranauna</i> R	<b>F</b> Striped Owl B <i>Pseudoscops clamator</i> R	<b>F</b> Turquoise-fronted Parrot B <i>Amazona aestiva</i> FDE
<b>III</b> Cinnamon Teal L <i>Anas georgica</i> R	<b>F</b> Cattle Egret L-VP <i>Bubulcus ibis</i> R	<b>I</b> Grey-necked Wood-Rail VP-T <i>Aramides cajaneus</i> R	<b>III</b> Brown-hooded Gull EA-C <i>Chroicocephalus maculipennis</i> RI	<b>F</b> Maroon-bellied Parakeet EA-B <i>Pyrrhura frontalis</i> FDE	
<b>F</b> Red Shoveler L <i>Anas cyanoptera</i> R	<b>II</b> Cooi Heron VP-C <i>Ardea cocoi</i> R	<b>I</b> Rufous-sided Crake VP <i>Laterallus melanophaius</i> R	<b>III</b> Grey-hooded Gull EA-C <i>Chroicocephalus cirrocephalus</i> RI	<b>I</b> Nacunda Hawk B <i>Chordeiles nacunda</i> R	
<b>I</b> Rosy-billed Pochard L <i>Netta peposaca</i> R	<b>I</b> Grea Egret EA-C <i>Ardea alba</i> R	<b>III</b> Red-and-white Crake VP <i>Laterallus leucopyrrhus</i> R	<b>F</b> Scissor-tailed Nighthawk B-P <i>Hydropsalis torquata</i> MA	<b>F</b> White-eyed Parakeet EA-B <i>Aratinga leucophthalma</i> FDE	
<b>III</b> Black-headed Swan L <i>Heteronetta atricapilla</i> R	<b>III</b> Whistling Heron VP <i>Syrigma sibilatrix</i> R	<b>III</b> Spotted Rail VP <i>Pardirallus maculatus</i> R	<b>F</b> Olrog's Gull EA-C <i>Larus atlanticus</i> RI	<b>I</b> Glittering-bellied Emerald A-B <i>Chlorostilbon lucidus</i> RV	
<b>III</b> Masked duck L <i>Nomonyx dominicus</i> R	<b>II</b> Snowy Egret L-VP <i>Egretta thula</i> R	<b>II</b> Plumbeous Rail VP <i>Pardirallus sanguinolentus</i> R	<b>II</b> Kelp Gull EA-C <i>Larus dominicanus</i> R	<b>I</b> Gilded Sapphire A-B <i>Hylocharis chrysura</i> RV	
<b>F</b> Lake Duck L <i>Oxyura vittata</i> R	<b>II</b> White-faced Ibis L <i>Plegadis chihi</i> R	<b>I</b> Common Gallinule L-VP <i>Gallinula cgaleata</i> R	<b>F</b> Snowy-crowned Tern EA-C <i>Sterna trudeaui</i> R	<b>I</b> Rufous-capped Antshrike A-B <i>Thamnophilus ruficapillus</i> R	
<b>I</b> White-tufted Grebe L <i>Rollandia rolland</i> R	<b>II</b> Bare-faced Ibis EA <i>Phimosus infuscatus</i> R	<b>I</b> Spot-flanked Gallinule L-VP <i>Gallinula melanops</i> R	<b>II</b> Picui Ground Dove B-T <i>Columbina picui</i> R	<b>III</b> Green Kingfisher L <i>Chloroceryle americana</i> R	
<b>N</b> Least Grebe L <i>Tachybaptus dominicus</i> FD	<b>III</b> Roseate Spoonbill EA <i>Platalea ajaja</i> RV	<b>N</b> Purple Gallinule VP <i>Porphyrio martinicus</i> R	<b>I</b> Rock Pigeon B-T <i>Columba livia</i> R	<b>I</b> Wren-like Rushbird VP <i>Phleocryptes melanops</i> R	
<b>I</b> Pied-billed Grebe L <i>Podilymbus podiceps</i> R	<b>II</b> Red-gartered Coot L-VP <i>Fulica armillata</i> R	<b>II</b> Red-fronted Coot L-VP <i>Fulica rufifrons</i> R	<b>I</b> Picazuro Pigeon B-T <i>Patagioenas picazuro</i> R	<b>III</b> White-barred Piculet B <i>Picumnus cirratus</i> R	
<b>II</b> Great Grebe C <i>Podiceps major</i> R	<b>F</b> White-tailed Kite EA-B <i>Elanus leucurus</i> R	<b>I</b> Red-fronted Coot L-VP <i>Fulica rufifrons</i> R	<b>III</b> Spot-winged Pigeon B <i>Patagioenas maculosa</i> R	<b>I</b> Checkered Woodpecker B <i>Veniliornis mixtus</i> R	


HABITAT: VP: Marsh Vegetation - EA: Air Space - A: Bushland - B: Woods - P: Grassland - T: Trails - C: Coast of river Plata. STATUS/SEASONAL DYNAMICS: R: Resident - RI: Winter Resident - RV: Summer Resident - MB: Summer Migrant (non-breeder) - MA: Summer Migrant (breeder) - MAP:

Winter Migrant (non-breeder) - FD: Out of range of distribution - FDE: ORD Established. DETECTABILITY: I: High - II: Medium - III: Low - F: Fortuitous - N: New

FIELD LIST OF RESERVA COSTANERA SUR - BIRDS RECORDED (May 2014-April 2015)

Cora Rimoldi, J. Simón Tagtachian and Germán Pugnali - www.reservacostanera.com.ar

Furnariidae (cont)		Tyrannidae (cont.)		Hirundinidae (cont.)		Thraupidae		Cardinalidae		Icteridae (cont.)	
<b>F</b> Sharp-billed Canastero	A-P	<b>II</b> Yellow-browed Tyrant	VP-P-A	<b>I</b> Blue-and-white Swallow	EA	<b>I</b> Red-crested Cardinal	T-A-B	<b>III</b> Hepatic Tanager	B	<b>II</b> Scarlet-headed Blackbird	VP
<i>Asthenes pyrrholeuca</i>	MAP	<i>Satrapa icterophrys</i>	R	<i>Pygochelidon cyanoleuca</i>	R	<i>Paroaria coronata</i>	R	<i>Piranga flava</i>	R	<i>Amblyramphus holosericeus</i>	R
<b>III</b> Sulphur-bearded Spinetail	P	<b>F</b> Dark-faced Ground-Tyrant	T	<b>I</b> Brown-chested Martin	EA-B	<b>I</b> Yellow-billed Cardinal	VP-T	<b>F</b> Glaucous-blue Grosbeak	B	<b>III</b> Unicolored Blackbird	VP
<i>Cranioleuca sulphurifera</i>	R	<i>Muscisaxicola maclovianus</i>	MAP	<i>Progne tapera</i>	MA	<i>Paroaria capitata</i>	R	<i>Cyanoloxia glaucocaerulea</i>	MA	<i>Agelasticus cyanopus</i>	R
<b>F</b> Yellow-chinned Spinetail	A	<b>F</b> Grey Monjita	VP	<b>I</b> Grey-breasted Martin	EA-B	<b>II</b> White-lined Tanager	B	<b>III</b> Ultramarine Grosbeak	A-B	<b>II</b> Yellow-winged Blackbird	VP
<i>Certhiaxis cinnamomeus</i>	R	<i>Xo9Imis cinereus</i>	MA	<i>Progne chalybea</i>	MA	<i>Tachyphonus rufus</i>	R	<i>Cyanocompsa brissonii</i>	R	<i>Agelasticus thilius</i>	R
<b>F</b> Chotoy Spinetail	A	<b>F</b> Black-backed Water-Tyrant	VP	<b>F</b> Southern Martin	EA-B	<b>F</b> Diademed Tanager	B	<b>III</b> Greyish Saltator	B	<b>III</b> Chestnut-capped Blackbird	VP
<i>Schoeniophylax phryganophilus</i>	R	<i>Fluvicola albiventer</i>	MA	<i>Progne elegans</i>	MA	<i>Stephanophorus diadematus</i>	R	<i>Saltator coerulescens</i>	R	<i>Chrysomus ruficapillus</i>	R
<b>III</b> Spix's Spinetail	A-B-P	<b>II</b> Cattle Tyrant	T-P	<b>I</b> White-rumped Swallow	EA-B	<b>F</b> Fawn-breasted Tanager	B	<b>II</b> Green-winged Saltator	B	<b>I</b> Bay-winged Cowbird	T-A-P
<i>Synallaxis spixi</i>	R	<i>Machetornis rixosa</i>	R	<i>Tachycineta leucorrhoa</i>	MA	<i>Pipraeidea melanonota</i>	RI	<i>Saltator similis</i>	R	<i>Agelaioides badius</i>	R
<b>I</b> Sooty-fronted Spinetail	A-B-P	<b>I</b> Great Kiskadee	VP-A-P	<b>F</b> Bank Swallow	EA-B	<b>I</b> Blue-and-yellow Tanager	B	<b>I</b> Golden-billed Saltator	A-B	<b>III</b> Screaming Cowbird	A-T
<i>Synallaxis frontalis</i>	R	<i>Pitangus sulphuratus</i>	R	<i>Riparia riparia</i>	MA	<i>Pipraeidea bonariensis</i>	R	<i>Saltator aurantirostris</i>	R	<i>Molothrus rufoaxillaris</i>	R
<b>Tyrannidae</b>		<b>I</b> Streaked Flycatcher	B	<b>F</b> Cliff Swallow	EA-B	<b>II</b> Sayaca Tanager	B	<b>Parulidae</b>		<b>III</b> Scrub Blackbird	A-B
<b>I</b> Small-billed Elaenia	B	<i>Myiodynastes maculatus</i>	RV	<b>F</b> Barn Swallow	EA	<i>Thraupis sayaca</i>	R	<b>II</b> Tropical Parula	B	<i>Dives warszewiczi</i>	FD
<i>Elaenia parvirostris</i>	MA	<b>F</b> Crowned Slaty Flycatcher	B	<b>F</b> Bank Swallow	EA	<i>Long-tailed Reed-Finch</i>	P-A	<i>Setophaga pitaiayumi</i>	R	<b>I</b> Shiny Cowbird	T-A
<b>F</b> Southern B.-Tyrannulet	B	<i>E.mpidonomus aurantioatrocristatus</i>	MA	<i>Hirundo rustica</i>	MA	<i>Donacospiza albifrons</i>	R	<i>Molothrus bonariensis</i>	R	<i>Molothrus bonariensis</i>	R
<i>Camptostoma obsoletum</i>	R	<b>I</b> Tropical Kingbird	B-P	<b>Troglodytidae</b>		<i>Black-and-rufous W.-Finch</i>	A-P	<i>Geothlypis aequinoctialis</i>	MA	<b>III</b> White-browed Blackbird	P
<b>F</b> Suiirí Flycatcher	B	<i>Tyrannus melancholicus</i>	MA	<b>I</b> House Wren	A-P-B	<i>Poospiza nigrorufa</i>	R	<i>Sturnella supercilialis</i>	MA	<i>Sturnella supercilialis</i>	MA
<i>Suiiriri suiiriri</i>	R	<b>I</b> Fork-tailed Flycatcher	B-P	<i>Troglodytes aedon</i>	R	<b>III</b> Grey-throated W.-Finch	B	<b>I</b> Golden-crowned Warbler	B	<b>Fringillidae</b>	
<b>F</b> Sooty Tyrannulet	VP	<i>Tyrannus savana</i>	MA	<b>Poliopitilidae</b>		<i>Poospiza cabanisi</i>	R	<i>Basileuterus culicivorus</i>	R	<b>I</b> Hooded Siskin	B-P-A
<i>Serpophaga nigricans</i>	R	<b>III</b> Swainson's Flycatcher	B	<i>Masked Gnatcatcher</i>	A-B	<i>Poospiza melanoleuca</i>	R	<b>Icteridae</b>		<i>Sporagra magellanica</i>	R
<b>I</b> White-crested Tyrannulet	A-B	<i>Myiarchus swainsoni</i>	MA	<i>Polioptila dumicola</i>	R	<b>I</b> Yellow Finch	A-P	<b>I</b> Solitary Black Caciue	B	<b>Passeridae</b>	
<i>Serpophaga subcristata</i>	R	<b>Cotingidae</b>		<b>III</b> White-tipped Plantcutter	A-B	<i>Sicalis flaveola</i>	R	<i>Cacicus solitarius</i>	R	<b>I</b> House Sparrow	T-A
<b>I</b> Straneck's Tyrannulet	A-B	<i>Phytotoma rutila</i>	MAP	<b>Turdidae</b>		<b>F</b> Grassland Yellow-Finch	A-P	<b>I</b> Variable Oriole	A-B	<i>Passer domesticus</i>	FDE
<i>Serpophaga griseicapilla</i>	MAP	<b>Tytiridae</b>		<b>I</b> Rufous-bellied Thrush	A-B-P-T	<i>Sicalis luteola</i>	R	<i>Icterus pyrrhopterus</i>	R		
<b>F</b> Bearded Tachuri	A.B	<b>F</b> White-naped Xenopsaris	B	<i>Turdus rufiventris</i>	R	<b>III</b> Great Pampa-Finch	P-A-T				
<i>Polystictus pectoralis</i>	MA	<i>Xenopsaris albinucha</i>	MA	<b>I</b> Creamy-bellied Thrush	A-B-T	<i>Embernagra platensis</i>	R				
<b>III</b> Warbling Doradito	VP-P	<b>F</b> Green-backed Becard	B	<i>Turdus amaurochalinus</i>	R	<b>F</b> Blue-black Grassquit	P-VP				
<i>Pseudocolaptes flaviventris</i>	R	<i>Pachyrhamphus viridis</i>	R	<b>Mimidae</b>		<i>Volatinia jacarina</i>	MA				
<b>F</b> Tawny-crowned P.-Tyrant	A.B	<b>I</b> White-winged Becard	B	<b>I</b> Chalk-browed Mockingbird	A-T	<b>F</b> Rusty-collared Seed-eater	P-VP				
<i>Euscarthmus meloryphus</i>	R	<i>Pachyrhamphus polychopterus</i>	MA	<i>Mimus saturninus</i>	R	<i>Sporophila collaris</i>	R				
<b>III</b> Many-coloured Rushbird	VP	<b>Vireonidae</b>		<b>I</b> White-banded Mockingbird	A-T	<b>I</b> Double-collared Seed-eater	P-A				
<i>Tachuris rubrigastra</i>	R	<b>F</b> Rufous-browed Peppershrike	B	<i>Mimus triurus</i>	MAP	<i>Sporophila caerulescens</i>	R				
<b>I</b> Bran-coloured Flycatcher	A-B	<i>Pyrocephalus rubinus</i>	MA	<b>Sturnidae</b>		<b>F</b> Red-crested Finch	A				
<i>Myiophobus fasciatus</i>	MA	<b>I</b> Red-eyed Vireo	B	<b>F</b> Crested Myna	B-VP	<i>Coryphospingus cucullatus</i>	RI				
<b>II</b> Vermilion flycatcher	A-B	<i>Cyclarhis gujanensis</i>	MA	<i>Acridotheres cristatellus</i>	FDE	<b>F</b> Yellow Cardinal	P-A				
<i>Pyrocephalus rubinus</i>	MA	<b>I</b> Spectacled Tyrant	A-B-T	<i>Sturnus vulgaris</i>	FDE	<i>Gubernatrix cristata</i>	R				
<b>F</b> Blue-billed Black-Tyrant	B	<i>Hymenops perspicillatus</i>	R	<b>Emberizidae</b>		<b>I</b> Rufous-collared Sparrow	T-P-A-B				
<i>Knipolegus cyanirostris</i>	RI					<i>Zonotrichia capensis</i>	R				



HABITAT: VP: Marsh Vegetation - EA: Air Space - A: Bushland - B: Woods - P: Grassland - T: Trails - C: Coast of river Plata. STATUS/SEASONAL DYNAMICS: R: Resident - RI: Winter Resident - RV: Summer Resident - MB: Summer Migrant (non-breeder) - MA: Summer Migrant (breeder) - MAP:

Winter Migrant (non-breeder) - FD: Out of range of distribution - FDE: ORD Established. DETECTABILITY: I: High - II: Medium - III: Low - F: Fortuitous - N: New

OBSERVATIONS: